

PCTWORLD INTELLECTUAL PROPERTY ORGANIZATION
International Bureau

INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(51) International Patent Classification ⁶: C07C 231/08	A1	(11) International Publication Number: WO 98/08805 (43) International Publication Date: 5 March 1998 (05.03.98)
(21) International Application Number: PCT/GB97/02335 (22) International Filing Date: 29 August 1997 (29.08.97) (30) Priority Data: 9618055.9 29 August 1996 (29.08.96) GB (71) Applicant (for all designated States except US): NYCOMED IMAGING AS [NO/NO]; Nycoveien 1-2, N-0401 Oslo (NO). (71) Applicant (for GB only): COCKBAIN, Julian, Roderick, Michaelson [GB/GB]; Frank B. Dehn & Co., 179 Queen Victoria Street, London EC4V 4EL (GB). (72) Inventors; and (75) Inventors/Applicants (for US only): GULBRANDSEN, Trygve [NO/NO]; Nycomed Imaging as, Nycoveien 1-2, N-0401 Oslo (NO). INGULDSTAD, Odd, Einar [NO/NO]; Nycomed Imaging as, Nycoveien 1-2, N-0401 Oslo (NO). HOLMAAS, Lars, Terje [NO/NO]; Nycomed Imaging as, Nycoveien 1-2, N-0401 Oslo (NO). (74) Agents: COCKBAIN, Julian et al.; Frank B. Dehn & Co., 179 Queen Victoria Street, London EC4V 4EL (GB).		(81) Designated States: AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, CA, CH, CN, CU, CZ, DE, DK, EE, ES, FI, GB, GE, GH, HU, IL, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MD, MG, MK, MN, MW, MX, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, UA, UG, US, UZ, VN, YU, ZW, ARIPO patent (GH, KE, LS, MW, SD, SZ, UG, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, ML, MR, NE, SN, TD, TG). Published <i>With international search report.</i>
(54) Title: PROCESS FOR THE PREPARATION OF CONTRAST AGENTS (57) Abstract The invention provides a process for the preparation of an N-alkyl-acylamino-phenyl-carboxylic acid or carboxylic acid derivative by liquid phase acylation and subsequent N-alkylation of a corresponding amino-phenyl-carboxylic acid (or carboxylic acid derivative), the improvement comprising the addition of an alkylating agent to a solution containing the reaction products of said acylation, to effect said N-alkylation.		

BEST AVAILABLE COPY

FOR THE PURPOSES OF INFORMATION ONLY

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AL	Albania	ES	Spain	LS	Lesotho	SI	Slovenia
AM	Armenia	FI	Finland	LT	Lithuania	SK	Slovakia
AT	Austria	FR	France	LU	Luxembourg	SN	Senegal
AU	Australia	GA	Gabon	LV	Latvia	SZ	Swaziland
AZ	Azerbaijan	GB	United Kingdom	MC	Monaco	TD	Chad
BA	Bosnia and Herzegovina	GE	Georgia	MD	Republic of Moldova	TG	Togo
BB	Barbados	GH	Ghana	MG	Madagascar	TJ	Tajikistan
BE	Belgium	GN	Guinea	MK	The former Yugoslav Republic of Macedonia	TM	Turkmenistan
BF	Burkina Faso	GR	Greece	ML	Mali	TR	Turkey
BG	Bulgaria	HU	Hungary	MN	Mongolia	TT	Trinidad and Tobago
BJ	Benin	IE	Ireland	MR	Mauritania	UA	Ukraine
BR	Brazil	IL	Israel	MW	Malawi	UG	Uganda
BY	Belarus	IS	Iceland	MX	Mexico	US	United States of America
CA	Canada	IT	Italy	NE	Niger	UZ	Uzbekistan
CF	Central African Republic	JP	Japan	NL	Netherlands	VN	Viet Nam
CG	Congo	KE	Kenya	NO	Norway	YU	Yugoslavia
CH	Switzerland	KG	Kyrgyzstan	NZ	New Zealand	ZW	Zimbabwe
CI	Côte d'Ivoire	KP	Democratic People's Republic of Korea	PL	Poland		
CM	Cameroon	KR	Republic of Korea	PT	Portugal		
CN	China	KZ	Kazakhstan	RO	Romania		
CU	Cuba	LC	Saint Lucia	RU	Russian Federation		
CZ	Czech Republic	LI	Liechtenstein	SD	Sudan		
DE	Germany	LK	Sri Lanka	SE	Sweden		
DK	Denmark	LR	Liberia	SG	Singapore		
EE	Estonia						

- 1 -

Process for the preparation of contrast agents

The invention relates to a process for the preparation of iodinated X-ray contrast agents, in particular non-ionic N-alkylated-acylamino-triiodophenyl compounds such as iohexol, iodixanol, iopentol and ioxilan.

Non-ionic iodinated X-ray contrast agents have achieved great commercial success over the past twenty years and accordingly improving the efficiency of their manufacture is of great importance.

The manufacture of non-ionic contrast agents includes production of the chemical drug substance (primary production), followed by formulation to drug product (secondary production). The drug substance is usually made in a multistep chemical synthesis, and is thoroughly purified before formulation. As with any commercial drug production it is important to optimize yield, process time and demand for expensive equipment. All these parameters depend both on the chemical reaction conditions and the work-up between each step. The number of steps in the overall synthesis will of course be of great importance and if the work-up between individual process steps could be omitted significant improvement in efficiency can be obtained, provided that sufficient quality and yield are maintained for the final product.

Omitting the work-up between two process steps means that the second step will be performed in the crude reaction solution resulting from the previous step. Potential major advantages includes:

- 2 -

- higher yield (through avoidance of loss of the intermediate in the work-up);
- elimination of isolation, purification, drying and analysis of the intermediate;
- significant reduction in equipment demand (eg. reactors, filters, driers, etc.); and
- significant reduction of the process time.

Despite these potential advantages, in practice intermediates are usually worked up between successive process steps. The reasons for this often include the facts that impurities may carry through the process and that the optimum solvent for the first reaction is usually not the same as for the second reaction.

The present multistage preparation of certain non-ionic X-ray contrast agents requires successive acylation and N-alkylation reactions. Thus for example in the production of iohexol (as described in SE-7706792-4 and by Gulbrandsen in Kjemi No. 6/90, pages 6-8), 5-amino-N,N'-bis(2,3-dihydroxypropyl)-2,4,6-triiodophthalamide (hereinafter 5-Amine) is acetylated to produce 5-acetamido-N,N'-bis(2,3-dihydroxypropyl)-2,4,6-triiodophthalamide (hereinafter 5-Acetamide) which is then N-alkylated to produce 5-[N-(2,3-dihydroxypropyl)-acetamido]-N,N'-bis(2,3-dihydroxypropyl)-2,4,6-triiodophthalamide (iohexol).

Similarly, in the production of iopentol as described in NO-160918, 5-Amine is acetylated to yield 5-Acetamide which is N-alkylated to produce 5-[N-(2-hydroxy-3-methoxypropyl)acetamido]-N,N'-bis(2,3-dihydroxypropyl)-2,4,6-triiodophthalamide (iopentol).

- 3 -

In the production of iodixanol described in NO-161368, 5-Amine is acetylated to yield 5-Acetamide which is then reacted with a bifunctional alkylating agent, a coupling agent, to yield 1,3-bis(acetamido)-N,N'-bis[3,5-bis(2,3-dihydroxypropylaminocarbonyl)-2,4,6-triiodophenyl]-2-hydroxypropane (iodixanol).

In the preparation of iohexol described above, the acylation is effected in acetic anhydride in the presence of a catalytic amount of sulphuric acid followed by concentration and addition of methanol. Thereafter water and sodium hydroxide (to pH 10-11) are added, base hydrolysis (to remove O-acyl groups) proceeds for 4-5 hours and 5-Acetamide is isolated from the reaction mixture by cooling and neutralization with hydrochloric acid. The precipitated 5-Acetamide is filtered, washed with water and dried. The 5-Acetamide is then dissolved in propylene glycol with the addition of sodium methoxide, the resulting methanol is stripped off and the N-alkylating agent (1-chloro-2,3-propanediol) is added. After the N-alkylation reaction is complete, the reaction mixture is evaporated to dryness and further purification steps (involving crystallization from a further solvent, butanol) are performed to yield the purified iohexol in a form suitable for use in secondary production.

The 5-Amine to 5-Acetamide acylation described for the preparation of iopentol in NO-160918 is similar, with the 5-Amine being acylated in acetic anhydride in the presence of a catalytic amount of p-toluene sulphonic acid. After cooling the reaction mixture, a precipitate forms which is filtered off and suspended in a mixture of methanol and water and hydrolysed under basic conditions (pH 11.5). The 5-Acetamide product is filtered off after cooling to ambient temperature, neutralization with HCl and further cooling to 3°C. The

- 4 -

5-Acetamide is washed with water and dried before being suspended in propylene glycol. Sodium hydroxide is added and when all solid material has dissolved, the N-alkylating agent (here 1-chloro-3-methoxy-2-propanol) is added. After the alkylation is quenched, several purification steps are again required before satisfactorily pure iopentol is obtained.

For the synthesis of iodixanol described in NO-161368, 5-Acetamide is prepared and worked up as described for iopentol above. It is then suspended in water and dissolved therein with the addition of sodium hydroxide. The N-alkylating agent, the coupling reagent epichlorohydrin, is then added. After the reaction is complete it is quenched with dilute hydrochloric acid and further purification steps are carried out in order to obtain satisfactorily pure iodixanol.

It has now been surprisingly found that the work-up of the 5-Acetamide before the subsequent N-alkylation may be avoided without unacceptable loss in yield or purity of the acylated, N-alkylated product and without undue complication of the purification procedure for that product.

Thus viewed from one aspect the invention provides a process for the preparation of an N-alkyl-acylamino-phenyl-carboxylic acid or carboxylic acid derivative by liquid phase acylation and subsequent N-alkylation of a corresponding amino-phenyl-carboxylic acid (or carboxylic acid derivative) characterised in that said N-alkylation is effected by addition of an alkylating agent to a solution containing the reaction products of said acylation, ie. those that remain in solution during the liquid phase acylation.

- 5 -

Viewed from an alternative aspect the invention provides a process for the preparation of an N-alkyl-acylamino-phenyl-carboxylic acid or carboxylic acid derivative compound comprising acylating an amino-phenyl-carboxylic acid (or carboxylic acid derivative) in a liquid phase, base hydrolysing the acylated product to remove O-acyl groups from the N-acyl-amino intermediate and, maintaining the liquid phase at a basic pH, N-alkylating the N-acylamino intermediate.

In the process of the invention, the aminophenyl carboxylic acid or carboxylic acid derivative starting product is preferably a compound having a total of three amino and carboxyl groups on the phenyl ring, eg. an aminoisophthalic acid or derivative or a diaminobenzoic acid or derivative. By carboxylic acid derivative is meant for example a salt, ester or amide, eg. an CONHR* or COOR* group where R* is optionally hydroxylated alkyl preferably optionally hydroxylated C₁₋₆ alkyl. Furthermore the starting compound is preferably a triiodophenyl compound, particularly an alkylamino-carbonyl-triiodophenyl compound and most particularly a 2,4,6-triiodo-2,5-bis(alkylamino-carbonyl)-aniline, such as 5-Amine for example. The alkyl moiety of any alkylaminocarbonyl group will preferably carry one or more hydroxyl groups and will typically contain up to 6, preferably up to 4 carbons.

The acylation of the starting compound may be effected using any convenient acylating agent, eg. an acetylating agent such as for example an acid halide or more preferably acetic anhydride.

Following the acylation, which is conveniently acid catalysed, the product is base hydrolysed, eg. at pH 11 to 12, to remove unwanted O-acyl groups without displacing the desired N-acyl groups. Base hydrolysis

- 6 -

may be with any organic or inorganic base but is preferably effected using an inorganic base, eg. an alkali metal hydroxide such as sodium hydroxide.

After the base hydrolysis, the solution pH is if necessary adjusted to an appropriate basic value for N-alkylation (generally within the range 10-12), and if desired further solvent (eg. a lower alcohol such as a C₁₋₄ alcohol, eg. methanol, optionally together with water) is added to the reaction mixture before, during or after addition to the reaction mixture of the alkylating agent, thereby ensuring that the non-isolated intermediate does not precipitate out.

Thus unlike prior successive acylation and N-alkylation reactions, the reaction medium is maintained at a basic pH and the N-acylated product (and the by-products) are not isolated from the reaction mixture before the subsequent N-alkylation step.

The N-alkylating agent used in the N-alkylation of the non-isolated intermediate may be any alkylating agent capable of introducing the desired alkyl group (or a precursor for a desired group) at the acylamino nitrogen. Where the acylated, N-alkylated product is intended to be a dimer (ie. containing two iodophenyl groups), the alkylating agent may be a bifunctional coupling agent.

For the preparation of iohexol, 1-halo-2,3-propanediols (eg. 1-chloro-2,3-propanediol) and glycidol are preferred alkylating agents. For the preparation of iopentol, 1-halo-3-methoxy-2-propanols (eg. 1-chloro-3-methoxy-propanol) are preferred. For the production of iodixanol a coupling agent such as a 1,3-halo-2-propanol or more preferably epichlorohydrin may be used.

- 7 -

The alkylation reaction may be terminated conventionally by quenching with acid, eg. HCl, and the acetylated, N-alkylated product may also be worked up conventionally, eg. by crystallization from an appropriate solvent or solvent mixture, eg. one or more C₁₋₆ alcohols.

Where the N-alkyl-acylamino product is a product such as iohexol, iopentol or iodixanol, this work-up yields a product which can be used for secondary production of an X-ray contrast medium. However for other end products, the product may be used as the starting material for further synthetic steps, eg. in the primary production of a non-ionic X-ray contrast agent.

Thus the process of the present invention allows N-acylation and N-alkylation to be effected as a one-pot synthesis achieving savings in equipment, time and materials without undue loss in yield and, surprisingly, achieving in the purification of the end product comparable or better purity levels to those achieved when the intermediate and the end product are both subject to work up stages.

The process of the present invention is particularly applicable to the preparation of the following N-alkyl-acylaminophenyl X-ray contrast agents: iomeprol, ioversol, ioxilan, iotrolan, ioxaglate, iodecimol, 2-iopyrol, 2-iopiperidol, iohexol, iopentol, iodixanol and the Schering products BP6 and BP8.

The invention will now be illustrated further by reference to the following non-limiting Examples.

- 8 -

EXAMPLE 1

5-Amino-N,N'-bis(2,3-dihydroxypropyl)-2,4,6-triiodoisophthalamide (388 g), acetic anhydride (838 g) and *p*-toluene sulfonic acid (2 g) were mixed and heated to reflux for 1.5 hours, before concentration to a highly viscous solution under reduced pressure. Methanol (44 g) was added to the solution in several portions with distillation under atmospheric pressure between each addition, followed by a final portion of methanol (266 g). A small amount of distilled water (40 ml) was added before stirring at 55°C overnight.

A portion (490 ml) of a solution made as described above was added to a jacketed reactor at about 30°C, and stirred with a speed of 150 rpm. The pH was raised by aqueous sodium hydroxide (50 w/v%, 190 ml) to about 12. 1-Chloro-2,3-propanediol (49 g) was added to the solution, and the pH was further adjusted by sodium hydroxide (50 w/v%, 25 ml) to about 11.5. Small amounts of additional 1-chloro-2,3-propanediol (total 6 ml) were added during the first 24 hours of the reaction. After a total reaction time of 99 hours, the solution was analyzed by reversed phase HPLC (water/acetonitrile as eluent) with the following results:

Iohexol	94.4%
5-Acetamide	1.9%
5-Amine	0.6%
Other impurities	3.1%

EXAMPLE 2

5-Amino-N,N'-bis(2,3-dihydroxypropyl)-2,4,6-triiodoisophthalamide (388 g), acetic anhydride (838 g) and *p*-toluene sulfonic acid (2 g) were mixed and heated to reflux for 1.5 hours, before concentration to a

- 9 -

highly viscous solution under reduced pressure. Methanol (44 g) was added to the solution in several portions with distillation under atmospheric pressure between each addition, followed by a final portion of methanol (266 g). A small amount of distilled water (40 ml) was added before stirring at 55°C overnight.

A portion (490 ml) of a solution made as described above was added to a jacketed reactor at about 35°C, and stirred with a speed of 150 rpm. The pH was raised by aqueous sodium hydroxide (50 w/v%, 200 ml) to about 12. Additional sodium hydroxide solution (10 ml) and 1-chloro-3-methoxy-2-propanol (42 g) were added, and the pH was adjusted back to about 12 by small amounts of sodium hydroxide solution. Further additions of 1-chloro-3-methoxy-2-propanol (total 21 g) were added after 2, 3, 4, 5 and 6 hours of reaction. After a total reaction time of 53 hours, the mixture was analyzed by HPLC (water/acetonitrile as eluent) with the following results:

Iopentol	91.4%
5-Acetamide	1.6%
5-Amine	0.3%
Other impurities	6.8%

- 10 -

Claims:

1. In a process for the preparation of an N-alkyl-acylamino-phenyl-carboxylic acid or carboxylic acid derivative by liquid phase acylation and subsequent N-alkylation of a corresponding amino-phenyl-carboxylic acid (or carboxylic acid derivative), the improvement comprising the addition of an alkylating agent to a solution containing the reaction products of said acylation, to effect said N-alkylation.
2. A process as claimed in claim 1 wherein said corresponding amino-phenyl carboxylic acid (or carboxylic acid derivative) is a compound having a total of three amino and carboxyl groups on the phenyl ring thereof.
3. A process as claimed in claim 1, wherein said corresponding amino-phenyl-carboxylic acid (or carboxylic acid derivative) is a triiodophenyl compound.
4. A process as claimed in claim 1, wherein said corresponding amino-phenyl-carboxylic acid (or carboxylic acid derivative) is an alkylamino-carbonyl-triiodophenyl compound.
5. A process as claimed in claim 1, wherein said corresponding amino-phenyl-carboxylic acid (or carboxylic acid derivative) is a 2,4,6-triiodo-2,5-bis(alkylamino-carbonyl)-aniline.
6. A process as claimed in claim 1, wherein said corresponding amino-phenyl-carboxylic acid (or carboxylic acid derivative) is 5-amino-N-N'-bis(2,3-dihydroxypropyl)-2,4,6-triiodophthalamide.

- 11 -

7. A process as claimed in claim 1, wherein said corresponding amino-phenyl-carboxylic acid (or carboxylic acid derivative) contains an alkylaminocarbonyl group carrying one or more hydroxyl groups and containing up to 6 carbon atoms.
8. A process as claimed in claim 1, wherein said acylation is effected using an acid halide or acetic anhydride.
9. A process as claimed in claim 1, wherein said alkylating agent is selected from the group consisting of 1-halo-2,3-propanediols, glycidol, 1-halo-3-methoxy-2-propanols, 1,3-halo-2-propanols and epichlorohydrin.
10. A process for the preparation of an N-alkyl-acylamino-phenyl-carboxylic acid or carboxylic acid derivative compound comprising acylating an amino-phenyl-carboxylic acid (or carboxylic acid derivative) in a liquid phase, base hydrolysing the acylated product to remove O-acyl groups from the resulting N-acyl-amino intermediate and, maintaining the liquid phase at a basic pH, N-alkylating said N-acylamino intermediate.
11. A process as claimed in claim 1 or claim 10, being a process for the preparation of a contrast agent selected from the group consisting of iomeprol, ioversol, ioxilan, iotrolan, ioxaglate, iodecimol, 2-iopyrol, 2-iopiperidol, iohexol, iopentol and iodixanol.

INTERNATIONAL SEARCH REPORT

International Application No
PCT/GB 97/02335

A. CLASSIFICATION OF SUBJECT MATTER
IPC 6 C07C231/08

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)
IPC 6 C07C

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practical, search terms used)

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
A	DE 27 26 196 A (NYEGAARD) 22 December 1977 cited in the application see claims 5-8	1,10
A	DE 22 16 627 A (LABORATOIRES ANDRE GUERBET) 19 October 1972 see claim 6 --- -/-	1,10

☒ Further documents are listed in the continuation of box C.

☒ Patent family members are listed in annex.

* Special categories of cited documents:

"A" document defining the general state of the art which is not considered to be of particular relevance

"E" earlier document but published on or after the international filing date

"L" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)

"O" document referring to an oral disclosure, use, exhibition or other means

"P" document published prior to the international filing date but later than the priority date claimed

"T" later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention

"X" document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone

"Y" document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art.

"Z" document member of the same patent family

Date of the actual completion of the international search

21 November 1997

Date of mailing of the international search report

10.12.97

Name and mailing address of the ISA

European Patent Office, P.B. 5818 Patentlaan 2
NL - 2280 HV Rijswijk
Tel. (+31-70) 340-2040, Tx. 31 651 epo nl,
Fax: (+31-70) 340-3016

Authorized officer

Voyiazoglou, D

INTERNATIONAL SEARCH REPORT

International Application No

PCT/GB 97/02335

C.(Continuation) DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
A	CHEMICAL ABSTRACTS, vol. 104, no. 24, 16 June 1986 Columbus, Ohio, US; abstract no. 213279a, JOSE L. MARTIN JIMENEZ ET AL: "Preparation of iodine-containing x-ray contrast agents" page 380; XP002047817 see abstract & ES 532 390 A (JUSTE S. A.) 16 June 1985 ---	1,10
A	EP 0 105 752 A (NYEGAARD) 18 April 1984 see claim 4 ---	1,10
A	WO 87 00757 A (COOK IMAGING) 12 February 1987 see claims 1,5 ---	1,10
A	WO 85 01727 A (BIOPHYSICA FOUNDATION) 25 April 1985 see claim 1 -----	1,10

INTERNATIONAL SEARCH REPORT

International application No.
PCT/GB 97/02335

Box I Observations where certain claims were found unsearchable (Continuation of Item 1 of first sheet)

This International Search Report has not been established in respect of certain claims under Article 17(2)(a) for the following reasons:

1. ☐ Claims Nos.:
because they relate to subject matter not required to be searched by this Authority, namely:
2. ☒ Claims Nos.: 1-4,7-11
because they relate to parts of the International Application that do not comply with the prescribed requirements to such an extent that no meaningful International Search can be carried out, specifically:
The claims are so broad that for determining the scope of a meaningful international Search due account has been taken of Rule 33.3 PCT; emphasis was put on claims 5,6 and on examples 1,2
3. ☐ Claims Nos.:
because they are dependent claims and are not drafted in accordance with the second and third sentences of Rule 6.4(a).

Box II Observations where unity of invention is lacking (Continuation of Item 2 of first sheet)

This International Searching Authority found multiple inventions in this international application, as follows:

1. ☐ As all required additional search fees were timely paid by the applicant, this International Search Report covers all searchable claims.
2. ☐ As all searchable claims could be searched without effort justifying an additional fee, this Authority did not invite payment of any additional fee.
3. ☐ As only some of the required additional search fees were timely paid by the applicant, this International Search Report covers only those claims for which fees were paid, specifically claims Nos.:
4. ☐ No required additional search fees were timely paid by the applicant. Consequently, this International Search Report is restricted to the invention first mentioned in the claims; it is covered by claims Nos.:

Remark on Protest

- ☐ The additional search fees were accompanied by the applicant's protest.
- ☐ No protest accompanied the payment of additional search fees.

INTERNATIONAL SEARCH REPORT

Information on patent family members

International Application No

PCT/GB 97/02335

Patent document cited in search report	Publication date	Patent family member(s)	Publication date
DE 2726196 A	22-12-77	GB 1548594 A	18-07-79
		BE 855580 A	12-12-77
		CH 630064 A	28-05-82
		DK 258477 A,B,	12-12-77
		FR 2354316 A	06-01-78
		HK 56683 A	25-11-83
		JP 1113258 C	16-09-82
		JP 53021137 A	27-02-78
		JP 56054310 B	24-12-81
		NL 7706385 A,C	13-12-77
		SE 441181 B	16-09-85
		SE 7706792 A	12-12-77
		US 4250113 A	10-02-81
DE 2216627 A	19-10-72	FR 2132530 A	24-11-72
		GB 1374918 A	20-11-74
EP 105752 A	18-04-84	DK 451983 A,B,	02-04-84
		HK 22891 A	04-04-91
		JP 2568166 B	25-12-96
		JP 59082355 A	12-05-84
		US 5204086 A	20-04-93
WO 8700757 A	12-02-87	AU 600672 B	23-08-90
		AU 6197486 A	05-03-87
		AU 650432 B	23-06-94
		AU 6686690 A	13-06-91
		AU 673286 B	31-10-96
		AU 7415594 A	05-01-95
		DE 3689330 D	05-01-94
		DE 3689330 T	19-05-94
		DK 178587 A	08-04-87
		EP 0233249 A	26-08-87
		JP 8013794 B	14-02-96
		JP 63500522 T	25-02-88
		NO 178106 B	16-10-95
		NO 178107 B	16-10-95
		US 4954348 A	04-09-90
		US 5035877 A	30-07-91

INTERNATIONAL SEARCH REPORT

Information on patent family members

International Application No

PCT/GB 97/02335

Patent document cited in search report	Publication date	Patent family member(s)	Publication date
WO 8501727 A	25-04-85	US 4584401 A	22-04-86
		AU 3550584 A	07-05-85
		EP 0160036 A	06-11-85
		JP 61500549 T	27-03-86
		US 5191119 A	02-03-93

**This Page is Inserted by IFW Indexing and Scanning
Operations and is not part of the Official Record**

BEST AVAILABLE IMAGES

Defective images within this document are accurate representations of the original documents submitted by the applicant.

Defects in the images include but are not limited to the items checked:

- ☐ **BLACK BORDERS**
- ☐ **IMAGE CUT OFF AT TOP, BOTTOM OR SIDES**
- ☐ **FADED TEXT OR DRAWING**
- ☐ **BLURRED OR ILLEGIBLE TEXT OR DRAWING**
- ☐ **SKEWED/SLANTED IMAGES**
- ☐ **COLOR OR BLACK AND WHITE PHOTOGRAPHS**
- ☐ **GRAY SCALE DOCUMENTS**
- ☐ **LINES OR MARKS ON ORIGINAL DOCUMENT**
- ☐ **REFERENCE(S) OR EXHIBIT(S) SUBMITTED ARE POOR QUALITY**
- ☐ **OTHER:** _____

IMAGES ARE BEST AVAILABLE COPY.

As rescanning these documents will not correct the image problems checked, please do not report these problems to the IFW Image Problem Mailbox.